



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

denser, components. The negative elements have, because of the relatively greater density which they inherited from an original state, been able to force the positive into higher positions during much of geologic time, so that they are characterized by a greater thickness of fairly continuous sediments. The positive elements found their levels of isostatic adjustment to be higher, so that they have more often constituted our land masses. The original relations of these components to one another has been greatly modified by tangential pressure and igneous intrusions, making occasional readjustments necessary; though the continent as a whole has maintained its equilibrium with the exceptionally heavy oceanic segments. H. H.

Peat, Essays on Its Origin, Uses, and Distribution in Michigan. By CHARLES A. DAVIS. A part of the report for 1906 of the Michigan Geological Survey, pp. 105-360, pls. XIII-XXXI, figs. 2-20. Lansing, 1907.

The author has very properly taken up the study of peat from a botanical standpoint, though geological factors are by no means ignored. Michigan peat is composed largely of the remains of plants which grew below or near the water-level; sphagnum forms only shallow superficial deposits. The living plant society varies as the development of a bog advances, for the changed conditions created by one group of species may enable an invading group to obtain a foothold where it was unable to live before the introduction of the earlier group. We cannot, therefore, expect the flora at present living in a swamp to indicate the quality of the underlying peat. A peculiar type of structureless peat was found which consisted largely of algal remains with occasional diatoms and an abundance of the three-celled pollen grains of conifers. Under proper conditions peat of this character would form a deposit like the structureless cannel coals of the Carboniferous. Many uses for which peat could be profitably employed are pointed out. In Michigan, especially, peat coke might be made to bear a close relationship to the iron industry. H. H.

The California Earthquake of 1906. By DAVID STARR JORDAN and Others. San Francisco: A. M. Robertson, 1907.

This volume contains a collection of articles, partly of a semi-popular nature, by D. S. Jordan, J. C. Branner, G. K. Gilbert, S. Tabor, F. Omori, H. W. Fairbanks, and Mary Austin, which had previously appeared in various publications, together with a new essay by Charles Derleth, Jr., on the effects produced by the earthquake on structures and structural